

PROJECTIONS IN GRAPH W^* -ALGEBRAS

ILWOO CHO

ABSTRACT. In this paper, we will consider the projections in a graph W^* -algebra $W^*(G)$. Let $T = L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ be a fixed operator in $W^*(G)$, where $w_1, \dots, w_n \in \mathbb{F}^+(G)$ and $u_{w_1}, \dots, u_{w_n} \in \{1, *\}$. We will find the conditions when this operator T is a projection. In other words, we will characterize the cases when $T = L_v$, for some $v \in V(G)$. In this paper, we show that there exists a vertex v such that $T = L_v$ if and only if the lattice path of T has the $*$ -axis-property if and only if $w_1^{t_1} \dots w_n^{t_n} = v$ in the free groupoid $\mathbb{F}(G)$, where $t_j = 1$ if $u_j = 1$ and $t_j = -1$ if $u_j = *$, for all $j = 1, \dots, n$.

In [2], Kribs and Power defined the free semigroupoid algebras and obtained some properties of them. Roughly speaking, graph W^* -algebras are W^* -topology closed version of free semigroupoid algebras. Throughout this paper, let G be a countable directed graph and let $\mathbb{F}^+(G)$ be the free semigroupoid of G , in the sense of Kribs and Power. i.e., it is a collection of all vertices of the graph G as units and all admissible finite paths, under the admissibility. As a set, the free semigroupoid $\mathbb{F}^+(G)$ can be decomposed by

$$\mathbb{F}^+(G) = V(G) \cup FP(G),$$

where $V(G)$ is the vertex set of the graph G and $FP(G)$ is the set of all admissible finite paths. Trivially the edge set $E(G)$ of the graph G is properly contained in $FP(G)$, since all edges of the graph can be regarded as finite paths with their length 1. We define a graph W^* -algebra of G by

$$W^*(G) \stackrel{def}{=} \overline{\mathbb{C}[\{L_w, L_w^* : w \in \mathbb{F}^+(G)\}]}^w,$$

where L_w and L_w^* are creation operators and annihilation operators of ξ_w on the generalized Fock space $H_G = l^2(\mathbb{F}^+(G))$ induced by the given graph G , respectively (See [1]). Let $w_1, \dots, w_n \in \mathbb{F}^+(G)$ and $u_1, \dots, u_n \in \{1, *\}$. Define

$$T = L_{w_1}^{u_1} \dots L_{w_n}^{u_n} \in W^*(G).$$

In this paper, we want to find the condition when T is a projection. i.e., we want to find the condition when there exists $v \in V(G)$ such that $T = L_v$. In [1], we already showed that $T = L_v$ if and only if the lattice path of T has the so-called the $*$ -axis-property with $*$ = v . In this paper, we will find one more condition which is

Key words and phrases. Graph W^* -Algebras. Shadowed Graph Algebras. Lattice Path Models.

equivalent to the $*$ -axis-property of T . We can show that $T = L_v$ is a projection if and only if the corresponding element $w_1^{t_1} \dots w_n^{t_n}$ is nothing but v in the free groupoid $\mathbb{F}(G)$, induced by the shadowed graph $G^\wedge = G \cup G^{-1}$ of the given graph G , with a certain relation on its free semigroupoid $\mathbb{F}^+(G^\wedge)$. Here, the shadowed graph $G^\wedge$ is the countable directed graph with

$$V(G^\wedge) = V(G) \text{ and } E(G^\wedge) = E(G) \cup E(G^{-1}),$$

where G^{-1} is the shadow of the graph G .

1. GRAPH W^* -ALGEBRAS

Let G be a countable directed graph and let $\mathbb{F}^+(G)$ be the free semigroupoid of G . i.e., the set $\mathbb{F}^+(G)$ is the collection of all vertices as units and all admissible finite paths of G . Let w be a finite path with its source $s(w) = x$ and its range $r(w) = y$, where $x, y \in V(G)$. Then sometimes we will denote w by $w = xwy$ to express the source and the range of w . We can define the graph Hilbert space H_G by the Hilbert space $l^2(\mathbb{F}^+(G))$ generated by the elements in the free semigroupoid $\mathbb{F}^+(G)$. i.e., this Hilbert space has its Hilbert basis $\mathcal{B} = \{\xi_w : w \in \mathbb{F}^+(G)\}$. Suppose that $w = e_1 \dots e_k \in FP(G)$ is a finite path with $e_1, \dots, e_k \in E(G)$. Then we can regard ξ_w as $\xi_{e_1} \otimes \dots \otimes \xi_{e_k}$. So, in [2], Kribs and Power called this graph Hilbert space the generalized Fock space. Throughout this paper, we will call H_G the graph Hilbert space to emphasize that this Hilbert space is induced by the graph.

Define the creation operator L_w , for $w \in \mathbb{F}^+(G)$, by the multiplication operator by ξ_w on H_G . Then the creation operator L on H_G satisfies that

$$(i) \quad L_w = L_{xwy} = L_x L_w L_y, \text{ for } w = xwy \text{ with } x, y \in V(G).$$

$$(ii) \quad L_{w_1} L_{w_2} = \begin{cases} L_{w_1 w_2} & \text{if } w_1 w_2 \in \mathbb{F}^+(G) \\ 0 & \text{if } w_1 w_2 \notin \mathbb{F}^+(G), \end{cases}$$

for all $w_1, w_2 \in \mathbb{F}^+(G)$.

Now, define the annihilation operator L_w^* , for $w \in \mathbb{F}^+(G)$ by

$$L_w^* \xi_{w'} \stackrel{def}{=} \begin{cases} \xi_h & \text{if } w' = wh \in \mathbb{F}^+(G) \\ 0 & \text{otherwise.} \end{cases}$$

The above definition is gotten by the following observation ;

$$\begin{aligned}
\langle L_w \xi_h, \xi_{wh} \rangle &= \langle \xi_{wh}, \xi_{wh} \rangle \\
&= 1 = \langle \xi_h, \xi_h \rangle \\
&= \langle \xi_h, L_w^* \xi_{wh} \rangle,
\end{aligned}$$

where $\langle, \rangle$ is the inner product on the graph Hilbert space H_G . Of course, in the above formula we need the admissibility of w and h in $\mathbb{F}^+(G)$. However, even though w and h are not admissible (i.e., $wh \notin \mathbb{F}^+(G)$), by the definition of L_w^* , we have that

$$\begin{aligned}
\langle L_w \xi_h, \xi_h \rangle &= \langle 0, \xi_h \rangle \\
&= 0 = \langle \xi_h, 0 \rangle \\
&= \langle \xi_h, L_w^* \xi_h \rangle.
\end{aligned}$$

Notice that the creation operator L and the annihilation operator L^* satisfy that

$$(1.1) \quad L_w^* L_w = L_y \quad \text{and} \quad L_w L_w^* = L_x, \quad \text{for all } w = xwy \in \mathbb{F}^+(G),$$

under the weak topology, where $x, y \in V(G)$. Remark that if we consider the von Neumann algebra $W^*(\{L_w\})$ generated by L_w and L_w^* in $B(H_G)$, then the projections L_y and L_x are Murray-von Neumann equivalent, because there exists a partial isometry L_w satisfying the relation (1.1). Indeed, if $w = xwy$ in $\mathbb{F}^+(G)$, with $x, y \in V(G)$, then under the weak topology we have that

$$(1.2) \quad L_w L_w^* L_w = L_w \quad \text{and} \quad L_w^* L_w L_w^* = L_w^*.$$

So, the creation operator L_w is a partial isometry in $W^*(\{L_w\})$ in $B(H_G)$. Assume now that $v \in V(G)$. Then we can regard v as $v = vvv$. So,

$$(1.3) \quad L_v^* L_v = L_v = L_v L_v^* = L_v^*.$$

This relation shows that L_v is a projection in $B(H_G)$ for all $v \in V(G)$.

Define the **graph W^* -algebra** $W^*(G)$ by

$$W^*(G) \stackrel{\text{def}}{=} \overline{\mathbb{C}[\{L_w, L_w^* : w \in \mathbb{F}^+(G)\}]}^w.$$

Then all generators are either partial isometries or projections, by (1.2) and (1.3). So, this graph W^* -algebra contains a rich structure, as a von Neumann algebra. (This construction can be the generalization of that of group von Neumann algebra.)

2. LATTICE PATH MODEL

Throughout this section, let G be a countable directed graph and let $W^*(G)$ be the graph W^* -algebra. Let $w_1, \dots, w_n \in \mathbb{F}^+(G)$ and let $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \in W^*(G)$ be an operator in $W^*(G)$. In this section, we will define a lattice path model for the random variable $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$. Recall that if $w = e_1 \dots e_k \in FP(G)$ with $e_1, \dots, e_k \in E(G)$, then we can define the length $|w|$ of w by k . i.e., the length $|w|$ of w is the cardinality k of the admissible edges $e_1, \dots, e_k$.

Definition 2.1. Let G be a countable directed graph and $\mathbb{F}^+(G)$, the free semi-groupoid. If $w \in \mathbb{F}^+(G)$, then L_w is the corresponding D_G -valued random variable in $(W^*(G), E)$. We define the lattice path l_w of L_w and the lattice path l_w^{-1} of L_w^* by the lattice paths satisfying that ;

- (i) the lattice path l_w starts from $*$ $= (0, 0)$ on the $\mathbb{R}^2$ -plane.
- (ii) if $w \in V(G)$, then l_w has its end point $(0, 1)$.
- (iii) if $w \in E(G)$, then l_w has its end point $(1, 1)$.
- (iv) if $w \in E(G)$, then l_w^{-1} has its end point $(-1, -1)$.
- (v) if $w \in FP(G)$ with $|w| = k$, then l_w has its end point (k, k) .
- (vi) if $w \in FP(G)$ with $|w| = k$, then l_w^{-1} has its end point $(-k, -k)$.

Assume that finite paths $w_1, \dots, w_s$ in $FP(G)$ satisfy that $w_1 \dots w_s \in FP(G)$. Define the lattice path $l_{w_1 \dots w_s}$ by the connected lattice path of the lattice paths $l_{w_1}, \dots, l_{w_s}$. i.e.e, l_{w_2} starts from $(k_{w_1}, k_{w_1}) \in \mathbb{R}^+$ and ends at $(k_{w_1} + k_{w_2}, k_{w_1} + k_{w_2})$, where $|w_1| = k_{w_1}$ and $|w_2| = k_{w_2}$. Similarly, we can define the lattice path $l_{w_1 \dots w_s}^{-1}$ as the connected path of $l_{w_s}^{-1}, l_{w_{s-1}}^{-1}, \dots, l_{w_1}^{-1}$.

Definition 2.2. Let G be a countable directed graph and assume that $L_{w_1}, \dots, L_{w_n}$ are generators of $W^*(G)$. Then we have the lattice paths $l_{w_1}, \dots, l_{w_n}$ of $L_{w_1}, \dots, L_{w_n}$, respectively in $\mathbb{R}^2$. Suppose that $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \neq 0_{D_G}$ in $W^*(G)$, where $u_{w_1}, \dots, u_{w_n} \in \{1, *\}$. Define the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ of nonzero $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ by the connected lattice path of $l_{w_1}^{t_{w_1}}, \dots, l_{w_n}^{t_{w_n}}$, where $t_{w_j} = 1$ if $u_{w_j} = 1$ and $t_{w_j} = -1$ if $u_{w_j} = *$. Assume that $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} = 0_{D_G}$. Then the empty set $\emptyset$ in $\mathbb{R}^2$ is the lattice path of it. We call it the empty lattice path. By LP_n , we will denote the set of all lattice paths of the D_G -valued random variables having their forms of $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$, including empty lattice path.

Also, we will define the following important property on the set of all lattice paths ;

Definition 2.3. Let $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \neq \emptyset$ be a lattice path of $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \neq 0_{D_G}$ in LP_n . If the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ starts from $*$ and ends on the $*$ -axis in $\mathbb{R}^+$, then

we say that the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ has the $*$ -axis-property. By LP_n^* , we will denote the set of all lattice paths having their forms of $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ which have the $*$ -axis-property. By little abuse of notation, sometimes, we will say that the D_G -valued random variable $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ satisfies the $*$ -axis-property if the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ of it has the $*$ -axis-property.

The following theorem shows that finding $E(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}})$ is checking the $*$ -axis-property of $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$.

Theorem 2.1. *Let $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \in W^*(G)$ be an operator, where $u_{w_1}, \dots, u_{w_n} \in \{1, *\}$. Then $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} = L_v$ if and only if $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ has the $*$ -axis-property.*

Proof. ($\Leftarrow$) Let $l = l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \in LP_n^*$. Suppose that $w_1 = vw_1v'_1$ and $w_n = v_nv_nv'_n$, for $v_1, v'_1, v_n, v'_n \in V(G)$. If l is in LP_n^* , then

$$(2.1) \quad \begin{cases} v_1 = v'_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = 1 \\ v_1 = v_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = * \\ v'_1 = v'_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = 1 \\ v'_1 = v_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = *. \end{cases}$$

By the definition of LP_n^* and by (2.1), if $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \in LP_n^*$, then there exists $v \in V(G)$ such that

$$(2.2) \quad \begin{cases} L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} = L_v, \\ \text{where} \quad \begin{cases} v = v_1 = v'_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = 1 \\ v = v_1 = v_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = * \\ v = v'_1 = v'_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = 1 \\ v = v'_1 = v_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = *. \end{cases} \end{cases}$$

This shows that $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} = L_v$.

($\Rightarrow$) Assume that $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} = L_v$, in $W^*(G)$. Let $l = l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \in LP_n$ be the lattice path of the D_G -valued random variable $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$. Trivially, $l \neq \emptyset$, since l should be the connected lattice path. Assume that this nonempty lattice path l is contained in $LP_n \setminus LP_n^*$. Then, under the same conditions of (2.1), we have that

$$(2.4) \quad \begin{cases} v_1 \neq v'_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = 1 \\ v_1 \neq v_n & \text{if } u_{w_1} = 1 \text{ and } u_{w_n} = * \\ v'_1 \neq v'_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = 1 \\ v'_1 \neq v_n & \text{if } u_{w_1} = * \text{ and } u_{w_n} = *. \end{cases}$$

Therefore, by (2.2), there is no vertex v satisfying $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} = L_v$. This contradicts our assumption. ■

3. SHADOWED GRAPH ALGEBRAS

In the previous chapter, we found one condition when the operator $T = L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ is a projection. In this chapter, we will find the equivalent concept. To do that we need to define the following new combinatorial object ;

Definition 3.1. Let G be a countable directed graph. Define the shadow G^{-1} of the graph G by a graph with

$$V(G^{-1}) = V(G) \quad \text{and} \quad E(G^{-1}) = \{e^{-1} : e \in E(G)\},$$

where e^{-1} is the reversely directed edge of e . Then we can define the shadowed graph $\hat{G} = G \cup G^{-1}$ of G by a graph with

$$V(\hat{G}) = V(G) \quad \text{and} \quad E(\hat{G}) = E(G) \cup E(G^{-1}).$$

Let $\mathbb{F}^+(\hat{G})$ be the free semigroupoid of the shadowed graph $\hat{G}$. Define the relation $\mathbf{R}$ on $\mathbb{F}^+(\hat{G})$ by

$$\mathbf{R} : \quad w^{-1}w = v' \quad \text{and} \quad ww^{-1} = v,$$

whenever $w = vw' \in FP(G)$, with $v, v' \in V(G)$. Define the free groupoid $\mathbb{F}(G)$ of G by the quotient set $\mathbb{F}^+(\hat{G}) / \mathbf{R}$.

Definition 3.2. Let G be a countable directed graph and let $\hat{G}$ be the shadowed graph of G . Also, let $\mathbb{F}(G)$ be the free groupoid of G . Let $l^2(\mathbb{F}(G))$ be the Hilbert space with its Hilbert basis $\{\xi_w : w \in \mathbb{F}(G)\}$. Then we can define the creation operator M_w of ξ_w , as usual, under the admissibility with the relation $\mathbf{R}$. Define a weak-closed algebra $\text{Alg}_w(G)$ by

$$\text{Alg}_w(G) \stackrel{\text{def}}{=} \overline{\mathbb{C}[\{M_w : w \in \mathbb{F}(G)\}]}^w.$$

This algebra $\text{Alg}_w(G)$ is called the shadowed graph algebra of G .

We will find the connection between our graph W^* -algebra $W^*(G)$ and the shadowed graph algebra $\text{Alg}_w(G)$. Notice that we can understand the algebra $\text{Alg}_w(G)$ as a weak-closed $*$ -algebra with the involution

$$M_w^* = M_{w^{-1}}, \quad \text{for all } w \in \mathbb{F}(G).$$

This admit us to conclude that

Theorem 3.1. *As weak-closed $*$ -algebras, the graph W^* -algebra $W^*(G)$ and the shadowed graph algebra $\text{Alg}_w(G)$ are isomorphic. $\square$*

The above theorem is easily proved because we can take the generator-preserving linear map,

$$L_w^{u_w} \in W^*(G) \longmapsto M_{w^{t_w}} \in \text{Alg}_w(G)$$

where

$$t_w = \begin{cases} 1 & \text{if } u_w = 1 \\ -1 & \text{if } u_w = *, \end{cases}$$

for all $w \in \mathbb{F}^+(G)$.

Let $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} \in W^*(G)$ be an operator, where $u_j \in \{1, *\}$. We can regard $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$ as $M_{w_1^{t_1}} \dots M_{w_n^{t_n}}$, by the previous theorem, where

$$t_j = \begin{cases} 1 & \text{if } u_j = 1 \\ -1 & \text{if } u_j = *, \end{cases}$$

for all $j = 1, \dots, n$, in $\text{Alg}_w(G)$. Notice that, by definition,

$$M_{w_1^{t_1}} \dots M_{w_n^{t_n}} = M_{w_1^{t_1} \dots w_n^{t_n}}$$

and

$$w_1^{t_1} \dots w_n^{t_n} \in \mathbb{F}(G) \quad \text{or} \quad w_1^{t_1} \dots w_n^{t_n} \notin \mathbb{F}(G).$$

By the previous discussion, we can have the following theorem ;

Theorem 3.2. *Let $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} \in W^*(G)$ be an operator, where $u_1, \dots, u_n \in \{1, *\}$, and let $M_{w_1^{t_1} \dots w_n^{t_n}} \in \text{Alg}_w(G)$ be the corresponding operator in $\text{Alg}_w(G)$, where $t_j = 1$ if $u_j = 1$ and $t_j = -1$ if $u_j = *$. Then $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} = L_v$, for some $v \in V(G)$, if and only if $w_1^{t_1} \dots w_n^{t_n} = v$.*

Proof. It is clear from the definition of the free groupoid. ■

Reference

- [1] I. Cho, Graph W^* -Probability Spaces, (2004), Preprint.
- [2] D.W. Kribs and S.C. Power, Free Semigroupoid Algebras, (2003), Preprint.

UNIV. OF IOWA, DEP. OF MATH, IOWA CITY, IA, U. S. A
E-mail address: `ilcho@math.uiowa.edu`